

THE BLACK SOLDIER FLY, *HERMETIA ILLUCENS*, AS A MANURE MANAGEMENT / RESOURCE RECOVERY TOOL

G. L. Newton, D. C. Sheppard, D. W. Watson, G. J. Burtle, C. R. Dove, J. K. Tomberlin, and E. E. Thelen

ABSTRACT

Insects play a significant role in recycling many forms of waste and other accumulated nutrients in the environment. Attempts to harness the power of insects for manure management have not been particularly successful or widely adopted. Although additional research is needed, the current state of the technology for utilizing the black soldier fly (*Hermetia*) for processing hen and swine manure appears to make it commercially feasible. *Hermetia* consume manure and convert the nutrients into larval insect mass, which contains 40+% protein and 30+% fat; thus a potential high protein, high energy animal feed. In the process, manure mass is reduced about 50%, with even greater reductions in N and P mass. Additional environmental benefits occur, and a recent analysis suggests that *Hermetia* manure management could increase net revenue by \$25,000! per year per layer house. For all applications, needed improved technology is associated with the adult life cycle and providing eggs throughout the year, and providing a warm environment for the larvae during winter. For conventional swine barn applications, economical methods for removal of excess liquid (urine) are also needed.

KEYWORDS. Hen manure, Swine manure Nutrient reduction, Bio-processing, Value-added, Insects

INTRODUCTION

Insects, especially various fly larvae (maggots) and some beetles, readily feed on fresh manure, converting residual protein and other nutrients into their biomass, which can be a high quality animal feedstuff. Considerable research has been conducted to understand this activity in order to exploit it for manure management. Lately, the emphasis has shifted from feedstuff production to potentially using insects to solve problems associated with the large amounts of manure produced at animal feeding operations (CAFOs). While incorporating and concentrating nutrients from manure into more valuable biomass (animal feedstuff), insect larvae reduce the nutrient concentration and bulk of the manure residue, thus reducing pollution potential 50-60% or more. As an added benefit, while occupying the manure they aerate and dry it, reducing odors. In addition, as a result of consuming and digesting microorganisms and the production of bacteriostatic, bactericidal, and/or fungicidal compounds in many species (Landi, 1960; Hoffmann and Hetru, 1992; Natori, 1995; Sherman et al., 2000), maggots modify the microflora of manure, potentially reducing harmful and undesirable species. This has been shown to occur with black soldier fly, potentially reducing harmful bacteria (Erickson, et al., 2004). Production of high-value feedstuff and reduction of the mass and pollution potential of the manure are the returns for good management of such a system.

While past research, and modest size demonstrations, have shown the potential of insect systems, they have not been adopted beyond some essentially subsistence situations. Three of the most important reasons for the lack of adoption include difficulties in adapting insect culture to modern animal production facilities, difficulties in producing eggs or larvae consistently on a year round basis, and effective, low cost methods for cold weather operation.

The black soldier fly (*Hermetia illucens*) (*Hermetia*) is a native insect common to the southeastern United States. Adults live and mate near larval habitat. *Hermetia* is not recognized as a pest

because the adult is not attracted to human habitation or foods (Furman et al. 1959). Adults do not need to eat, surviving on the large fat body stored from the larval stage. The larva of *Hermetia* is a voracious consumer of decaying organic matter including kitchen waste, spoiled feed, and manure. They flourished in open-sided caged layer houses that, in the past, were common in the southern U.S. In these situations soldier fly larvae were present by the millions, but only a few ovipositing females were observed. These unmanaged populations eliminated house fly breeding and reduced manure residue, but feedstuff harvest was never attempted. Prepupae harvest from managed *Hermetia* populations could be a viable addition to confined livestock enterprises.

HERMETIA MANURE MANAGEMENT SYSTEMS

The simplest *Hermetia* manure management system utilizes wild fly populations to digest the manure. The larvae occur in very dense populations on organic wastes as diverse as manures, coffee bean pulp, vegetables, catsup and carrion. Mature larvae (prepupae) can be harvested and used as a feedstuff. The *Hermetia* system developed by Sheppard et al. (1994) converted poultry manure to a 42% protein, 35% fat feedstuff at an 8% dry matter conversion rate (Sheppard et al., 1994). This system also controlled house flies (Sheppard, 1983) and reduced manure volume by 50% (Sheppard 1983), including a 24% reduction in total nitrogen concentration (62% reduction of N mass) (Sheppard et al., 1998).

Hermetia are most easily managed directly under caged layers (or other animals) in concrete basins. No separate facility or special equipment is needed for production or harvest. This is possible because of the migratory habits of the prepupae. This migration occurs because larvae need to leave the manure to successfully pupate to an adult. At this stage they are at their maximum size, with a large store of fat to sustain them through metamorphosis. Migrating prepupae have evacuated their digestive tract and no longer feed, but use their mouthparts to pull their body along in the quest for a safe pupation site. Pupation takes a minimum of 10 days so collections can be stockpiled prior to processing or utilization. In a 5-month *Hermetia* season a 100,000 bird caged layer house could produce 48,000 kg of prepupae suitable for feed. Many of the difficulties associated with utilization of *Hermetia* for manure processing in enclosed animal facilities, and for other animals, including swine and dairy calves, have been resolved, although additional improvements would allow widespread adoption of the system.

Swine System One

Recent developments in rearing *Hermetia* allow for waste management in fully enclosed buildings (Figure 1). Swine waste collected by conveyor belt was separated into manure solids and urine plus excess water. Collected manure solids were delivered to the larval culture basin. The larval culture basin contained 85,000 to 100,000 mixed aged larvae/m². A 35° ramp along opposing walls of the manure pit directed the migrating prepupae to a gutter at the top. This gutter directed prepupae to collection containers. A portion of the prepupae were saved and used to support the adult soldier fly colony. Eggs from the adult colony were used to maintain larval densities sufficient to digest the manure. The remaining prepupae were dried and processed for rendering or feed preparation. The digested swine manure could undergo further treatment or be land applied as a soil amendment.

Experimental Results

Eighteen pigs were housed over a manure collection belt during the grow-finish period. Manure was removed once per day and distributed over a culture basin containing mixed age *Hermetia* larvae. Over the course of the trial, manure mass total solids was reduced by 56% compared to fresh manure. The composition of the self-harvested prepupae is shown in Table 1, and the composition of the residue remaining after larval culture, compared to the sampled fresh manure is shown in Table 2.




Figure 0. *Hermetia* larvae production away from pigs

Table 1. Essential amino acid, mineral, and nutrient content of *Hermetia* prepupae fed on swine manure.

Essential Amino Acids (%)		Mineral and Other	
Methionine	0.83	P	0.88%
Lysine	2.21	K	1.16%
Leucine	2.61	Ca	5.36%
Isoleucine	1.51	Mg	0.44%
Histidine	0.96	Mn	348 ppm
Phenylalanine	1.49	Fe	776 ppm
Valine	2.23	Zn	271 ppm
l-Arginine	1.77	Crude protein	43.2%
Threonine	1.41	Ether extract	28.0%
Tryptophan	0.59	Ash	16.6%

Table 2. Composition of fresh pig manure compared to the residue remaining after *Hermetia* culture.

Element	Pig manure		Hermetia residue		Change %
	ppm	SEM	ppm	SEM	
N	923.7	44.4**	414.52	6.17	-55.1
P	676.2	37.9**	378	13.1	-44.1
K	358.7	19.8**	169.34	7.07	-52.8
Ca	969.3	62.5**	425	19.4	-56.2
Mg	299.3	16.9**	175.96	7.08	-41.2
S	80.31	4.33**	44.44	1.38	-44.7
Fe	6.63	0.31 ^{ns}	6.8	0.54	+2.6
Mn	12.8	0.75**	6.02	0.23	-53.0
Zn	23.53	1.09**	12.91	0.39	-45.1
Cu	14.85	1.45**	8.05	0.32	-45.8
B	0.32	0.02**	0.16	0	-50.0
C	11,248	497**	4,232.6	36.8	-62.4
Na	99.93	5.59**	48.15	2.04	-51.8

** = significant at $P < 0.001$, One way ANOVA, Minitab 1997; ns = not significant.

The results presented in Table 1 illustrate the potential for *Hermetia* prepupae to be used as a feed ingredient for most domestic animals. The composition of the prepupae was similar to that in earlier trials, except that the lipid content was somewhat lower (28% compared to 32-36%).

The comparison of fresh manure with the residue remaining after larval culture (Table 2) shows that the concentrations of N and minerals, with the exception of iron, were drastically reduced by the feeding larvae. If the 56% reduction in the mass of the total solids is taken into account, total mass reductions for many of the elements shown in Table 2 were near 80%. Compared with fresh manure, much less crop land area would be needed to manage the residue, compared to manure.

Swine System Two

If urine and spilled water can be drained or diverted, *Hermetia* larvae can be cultured directly beneath pigs housed on slatted floors. Such a system would appear to be readily adaptable to pigs housed in high-rise buildings. A combination of slotted and screened standpipes and laterals within the basin (based on the Ozyboyd Vertical Drainage Bed) has been found to be acceptable for excess liquid removal. These drains are attached to pipes beneath the floor, and can be

removed for basin cleaning. The second system for culture on swine manure is illustrated in Figure 2.




Figure 2. Culture of *Hermetia* larvae directly beneath pigs.

Experimental Results

During summer 2004, two manure basin liquid removal systems were tested. Both systems were tested with two pens of six pigs each (initial wt., 21 kg), housed over the individual basins. Treatment 1 consisted of 16 slotted and screened standpipes plus 4 slotted and screened laterals while treatment 2 was a double system with two sets of stand pipes and laterals with discharges that were alternately opened or closed at 14 day intervals. Alternately opening and closing the drainage system actually resulted in more "blinding" and restricted flow than leaving the drainage system open continuously.

During the grow-finish period, larvae collections were 0.214 kg/pig/day for treatment 1 and 0.153 kg/pig/day ($P < .03$) for treatment 2 (including the first 4-5 weeks when the larvae population was being established and essentially no prepupae were collected). Equivalent prepupal collections would equal to 64,000 kg/year for a 1,000 head finishing house having 2.5 turns of pigs/year.

Layer Hen System

Much of the interest in the use of the black soldier fly for manure processing and house fly control was prompted by observations of natural populations in old-style, open sided layer houses. The evolution to newer type housing for laying hens eliminated most of the wild *Hermetia* population within hen houses. However, it should be possible to introduce controlled populations of *Hermetia* to high-rise hen houses in order to process the manure into a lower nutrient, soil amendment and a high value feedstuff. This has been demonstrated in a small pilot scale house and proposed designs have been developed for full scale houses. The general design for such a house is shown in Figure 3.





Figure 3. Hen house with *Hermetia* manure processing.



REFERENCES

1. Erickson, M.C., M. Islam, C. Sheppard, J. Liao and M.P. Doyle. 2004. Reduction of *Escherichia coli* 0157:H7 and *Salmonella enterica* Serovar Enteritidis in chicken manure by larvae of the black soldier fly. *J. Food Protection*. 67: 685-690.
2. Furman, D. P., R. D. Young. & E. P. Catts. 1959. *Hermetia illucens* (Linnaeus) as a factor in the natural control of *Musca domestica* Linnaeus. *J. Econ. Entomol.*, 52: 917-921.
3. Landi, S. 1960. Bacteriostatic effect of hemolymph of larvae of various botflies. *Canadian J. Microbiol.*, 6: 115-119.
4. Natori, S. 1995. Antimicrobial proteins of insects and their clinical application. *Nippon Rinsho*. 5:1297-1304.
5. Sheppard, Craig. 1983. House fly and lesser house fly control utilizing the Black Soldier Fly in manure management systems for caged laying hens. *Environ. Entomol.* 12: 1439-1442.
6. Sheppard, D. C., G. L. Newton, S. Thompson, J. Davis, G. Gascho, K. Bramwell. 1998. Using soldier flies as a manure management tool for volume reduction, house fly control and reduction, house fly control and feedstuff production (pp 51-52). (In) Sustainable Agriculture Research and Education, Southern Region, 1998 Annual Report. Gwen Roland, Editor.
7. Sheppard, D. C., G. L. Newton, S. A. Thompson and S. E. Savage. 1994. A value added manure management system using the Black Soldier Fly. *Bioresource Technology*. 50: 275-279.
8. Sherman, R.A., M.J.R. Hall, S. Thomas. 2000. Medicinal maggots: An ancient remedy for some contemporary afflictions. *Ann. Rev. Entomol.* 45:55-81.